

TEBUCONAZOLE	GROUP	3	FUNGICIDE
AZOXYSTROBIN	GROUP	11	FUNGICIDE



HELMSTAR PLUS SC

FIFRA 2(ee) Recommendation
EPA Reg. No. 74530-69
FOR DISTRIBUTION AND USE ONLY IN THE FOLLOWING STATES: GEORGIA, IOWA, ILLINOIS, NEBRASKA, MINNESOTA, INDIANA, KANSAS, KENTUCKY, OHIO, MISSOURI, PENNSYLVANIA, SOUTH DAKOTA AND WISCONSIN
THIS RECOMMENDATION, WHICH CONTAINS ADDITIONAL DIRECTIONS FOR USE, IS MADE AS PERMITTED UNDER FIFRA SECTION 2(ee) AND HAS NOT BEEN SUBMITTED TO OR APPROVED BY THE EPA.

This recommendation for HELMSTAR PLUS SC herbicide is valid until December 31, 2028 or until withdrawn, cancelled, or suspended, whichever comes first.

HELMSTAR PLUS SC TO AID IN SUPPRESSION OF TAR SPOT IN CORN

IT IS A VIOLATION OF FEDERAL LAW TO USE THIS PRODUCT IN A MANNER INCONSISTENT WITH ITS LABELING. ALL APPLICABLE DIRECTIONS, RESTRICTIONS AND PRECAUTIONS ON THE EPA REGISTERED LABEL MUST BE FOLLOW.

THESE USE DIRECTIONS MUST BE IN THE POSSESSION OF THE USER AT THE TIME OF PESTICIDE APPLICATION.

CROP	PEST - AIDS IN SUPPRESSION	PRODUCT RATE PER APPLICATION (FL OZ/A)	MAXIMUM PRODUCT RATE PER YEAR (FL OZ/A)	MINIMUM TIME FROM APPLICATION TO HARVEST – DAYS (PHI)
Corn Field, Popcorn, and Seed Sweet corn	Tar Spot (<i>Phyllachora maydis</i>)	10.8	43.2	<p align="center">Field and Pop Corn</p> <p align="center">21 days for forage 36 days for grain or fodder.</p> <p align="center">Sweet Corn</p> <p align="center">7 days of harvest for ears or forage</p> <p align="center">49 days before the harvest of fodder.</p>

Apply **HELMSTAR PLUS SC** in a protective spray schedule or when weather conditions are favorable for disease development. Repeat applications at 7- to 14-day intervals, or as necessary to maintain suppression. Shorten the interval under heavy disease pressure.

Restrictions for Use of Adjuvants or Crop Oil in Corn.

DO NOT use adjuvants or crop oil after the V8 stage and prior to the VT stage. (The VT stage is defined as when the last branch of the tassel is completely visible outside of the whorl).

A compatibility agent, another fungicide, or an insecticide may be included in the tank mix, if needed, and labeled for use on corn. Refer to the adjuvant and other tank mix pesticide product labels for specific use directions and restrictions.

Always follow the most restrictive label.

Consult a HELM Agro representative or local agricultural authority for more information concerning additives.

For best results, tank mix **HELMSTAR PLUS SC** with the lowest labeled rate of a spray adjuvant (non-ionic surfactant, crop oil concentrate, or blend) at the manufacturers recommended rates to obtain sufficient coverage. Adjuvants that contain some form of silicone can contribute to phytotoxicity.

Use a higher water volume for aerial application if equipment and/or conditions will not provide good coverage.

RESTRICTIONS:

- Do not apply more than 43.2 fl oz/A/season of HELMSTAR PLUS SC per crop.
- Do not apply more than 0.675 lb. a.i. Tebuconazole containing products/A/year.
- Do not apply more than 2.0 lbs. a.i. Azoxystrobin containing products/A/year.
- Excluding sweet corn, restricted-entry interval (REI) = 12 hours.
- For sweet corn, restricted-entry interval (REI) = 19 days.

ALL APPLICABLE DIRECTIONS, RESTRICTIONS, WORKER PROTECTION STANDARDS REQUIREMENTS AND PRECAUTIONS ON THE EPA REGISTERED HELMSTAR PLUS SC LABEL MUST BE FOLLOWED. THIS 2(ee) RECOMMENDATION MUST BE IN THE POSSESSION OF THE USER AT THE TIME OF PESTICIDE APPLICATION.

(RV061724)

Manufactured For:
HELM Agro US, Inc.
 401 E. Jackson St., Suite 1600
 Tampa, FL 33602
 Phone: 813.621.8846
info@hmagro.com