



FIFRA 2(ee) Recommendation
EPA Reg. No. 71512-37-74530
FOR DISTRIBUTION AND USE ONLY IN THE FOLLOWING STATES: NORTH CAROLINA, SOUTH CAROLINA and VIRGINIA
THIS RECOMMENDATION, WHICH CONTAINS ADDITIONAL DIRECTIONS FOR USE, IS MADE AS PERMITTED UNDER FIFRA SECTION 2(ee) AND HAS NOT BEEN SUBMITTED TO OR APPROVED BY THE EPA.

This recommendation for Reviton herbicide is valid until December 31, 2026 or until withdrawn, cancelled, or suspended, whichever comes first.

FOR CONTROL OF FIELD PANSY IN PREPLANT BURNDOWN

IT IS A VIOLATION OF FEDERAL LAW TO USE THIS PRODUCT IN A MANNER INCONSISTENT WITH ITS LABELING. ALL APPLICABLE DIRECTIONS, RESTRICTIONS AND PRECAUTIONS ON THE EPA REGISTERED LABEL MUST BE FOLLOWED.

THESE USE DIRECTIONS MUST BE IN THE POSSESSION OF THE USER AT THE TIME OF PESTICIDE APPLICATION.

CROP	PEST	REVITON RATE OF APPLICATION
None	Field Pansy <i>(Viola bicolor)</i>	1 -2 fl oz/A

DIRECTION FOR USE:

- Always include an MSO at 1% v/v.
- Tank mix with Glyphosate at specified label rate on label, but not less than 1.125 lb. ae/A.
- Aerial applications should be applied with a minimum of 3 to 5 gallons per acre.
- Ground applications should be applied with a minimum of 10 to 15 gallons per acre.

RESTRICTIONS:

- Do not apply Reviton at a rate greater than 2 fl oz (0.045 lbs. ai) per acre. Do not exceed 6 fl oz (0.134 lbs. ai) per acre per year.
- Follow all plantback intervals listed on the Reviton label.
- Do not allow livestock to graze for 21 days after application.

ALL APPLICABLE DIRECTIONS, RESTRICTIONS, WORKER PROTECTION STANDARDS REQUIREMENTS AND PRECAUTIONS ON THE EPA REGISTERED REVITON LABEL MUST BE FOLLOWED.

THIS 2(ee) RECOMMENDATION MUST BE IN THE POSSESSION OF THE USER AT THE TIME OF PESTICIDE APPLICATION.

(RV050223)

HELM Agro US, Inc.
401 E. Jackson St., Suite 1400
Tampa, FL 33602
Phone: 813.621.8846
Fax: 813.621.0763
info@helmagro.com helmagro.com