

FIFRA Section 24(c) Special Local Need Label
FOR USE AND DISTRIBUTION ONLY WITHIN THE STATE OF ARKANSAS

| | | | |
|------------|-------|----|-----------|
| TIAFENACIL | GROUP | 14 | HERBICIDE |
|------------|-------|----|-----------|

REVITON

A nonselective burndown herbicide with the active ingredient TERGEO™

Aerial Application in the State of Arkansas

Active Ingredient

| | |
|-------------------|--------|
| Tiafenacil* | 30.0% |
| Other Ingredients | 70.0% |
| Total | 100.0% |

*methyl N-[2-[[2-chloro-5-[3,6-dihydro-3-methyl-2,6-dioxo-4-(trifluoromethyl)-1(2H)-pyrimidinyl]-4-fluorophenyl]thio]-1-oxopropyl]-β-alaninate

Reviton is formulated as a suspension concentrate (SC) and contains 2.83 pounds of active ingredient per gallon of formulated product (339 grams per liter).

EPA Reg. No. 71512-37-74530

Expiration: December 31, 2025

SLN No. AR22-0001

KEEP OUT OF REACH OF CHILDREN CAUTION/PRECAUCION

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.
(If you do not understand this label, find someone to explain it to you in detail.)

This label is valid until December 31, 2025 or until otherwise amended, withdrawn, cancelled or suspended.

SEE LABEL BOOKLET FOR FIRST AID, ADDITIONAL PRECAUTIONARY STATEMENTS AND DIRECTIONS FOR USE INCLUDING STORAGE AND DISPOSAL

READ ENTIRE LABEL CAREFULLY AND USE ONLY AS DIRECTED.

DIRECTIONS FOR USE

IT IS A VIOLATION OF FEDERAL LAW TO USE THIS PRODUCT IN ANY MANNER INCONSISTENT WITH ITS LABELING.

This label must be in the possession of the user at the time of application.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

ALL APPLICABLE DIRECTIONS, RESTRICTIONS AND PRECAUTIONS ON THE PRODUCT LABEL BOOKLET MUST BE FOLLOWED, INCLUDING STATEMENTS PERTAINING TO THE WORKER PROTECTION STANDARDS, ON THE EPA REGISTERED LABEL. BEFORE USING REVITON AS PERMITTED ACCORDING TO THIS SUPPLEMENTAL LABEL, READ AND FOLLOW ALL APPLICABLE DIRECTIONS, RESTRICTIONS AND PRECAUTIONS ON THE EPA REGISTERED LABEL AND THE ATTACHED PESTICIDE PRODUCT CONTAINER.

PRODUCT USES & APPLICATION INSTRUCTIONS:

This SLN Label for **Reviton** is registered for:

- 1) preplant and preemergence burndown use in corn (all types, except sweet corn and popcorn), and
- 2) preplant use in soybean and cotton.

Restrictions

- DO NOT apply this product to residential areas.
- DO NOT apply this product through any type of irrigation system.
- For any combination of cropping systems, DO NOT apply more than a maximum cumulative amount of 0.223 lb ai per acre per year.
- A 25 foot buffer for ground applications and a 150 foot buffer for aerial applications must be maintained between the point of direct application and the closest downwind edge of sensitive terrestrial habitats (including grasslands, forested areas, shelter belts, woodlots, hedgerows, riparian areas, shrub lands, crop lands), and estuarine/marine habitats.

Spray Volume – Aerial Application:

The minimum spray volume for aerial applications of **Reviton** is 3 or more gallons of final spray solution per acre. Adequate spray coverage is essential for optimal weed control. When targeting dense weed populations and/or larger weeds, use a minimum of 5 gallons of final spray solution per acre.

Mandatory Spray Drift Management

- Applicators must not spray during temperature inversions.
- For aerial applications, do not apply when wind speeds exceed 10 mph at the application site. The boom length must be 75% or less of the wingspan for fixed-wing aircraft and 90% or less of the rotor diameter for helicopters.
- For aerial applications, applicators must use ½ swath displacement upwind at the downwind edge of the field.
- For aerial applications, the release height must be no higher than 10 feet from the top of the crop canopy or ground, unless a greater application height is required for pilot safety.
- For ground boom applications, apply with the release height no more than 3 feet above the ground or crop canopy.
- For ground boom-less application (e.g., roadsides, rights of ways) there will be no mandatory release height restrictions due to equipment having physical limitations for achieving a height of 3 feet. There will be advisory spray drift language informing applicators to set nozzles at the lowest effective height to help reduce the potential for spray drift. The droplet size for these applications are also typically coarser, thus reducing the risk of spray drift.
- For ground applications, do not apply when wind speeds exceed 10 miles per hour at the application site.
- Applicators are to select a nozzle and pressure combination that delivers a medium or coarser droplet size (ASABE S572 for ground applications and ASABE S641 for aerial applications).

Adjuvants:

For best results, use a methylated seed oil (MSO) when applying **Reviton**, or reduced performance can occur. When using an MSO, always use a product that contains at least 80% high quality petroleum (mineral). MSO should be applied at a concentration equal to 1% v/v (1 gallon per 100 gallons spray carrier) of the final spray volume.

If using a crop oil concentrate (COC), always use a COC with modified vegetable oil with at least 15% surfactant emulsifier.

If a nonionic surfactant (NIS) is used, reduced performance can occur. If using an NIS, always use NIS containing at least 60% NIS, at a concentration equal to 0.25% v/v (2 pints per 100 gallons spray volume) of the final spray volume.

The addition of an ammonium nitrogen fertilizer, either a 28% or 32% N urea ammonium nitrate (UAN) or a spray grade ammonium sulfate (AMS), to the final spray solution is allowed. If UAN or AMS is added to the spray mixture, add UAN at a concentration of 2.5% v/v (2.5 gallons per 100 gallons or spray volume) and add AMS at a concentration of 8.5 lbs product per 100 gallons of the final spray volume.

CROP – SPECIFIC USE INSTRUCTIONS**Corn (All Types, Except Sweet Corn and Popcorn)**

The maximum single application rate is 3 fl oz per acre (0.067 lb ai per acre). Do not exceed 6 fl oz per acre per year (0.134 lb ai per acre per year).

| Application Timing | Rate Range (fl oz/A) | Additional Information & Restrictions |
|---------------------------|-----------------------------|---|
| Preplant Burndown | 1 to 3 | <ul style="list-style-type: none"> • Apply as a broadcast spray using conventional low-pressure ground spray equipment or by aerial application. • Follow manufacture's recommendations for spraying pressure. • Review and follow restrictions from the spray drift management section. • Do not reapply within 14 days. • Use higher rate for dense and/or mature weed infestations. |
| Preemergence Burndown | 1 to 3 | <ul style="list-style-type: none"> • Apply as a broadcast spray using conventional low-pressure ground spray equipment or by aerial application. • Follow manufacture's recommendations for spraying pressure. • Review and follow restrictions from the spray drift management section. • Do not reapply within 14 days. • Use higher rate for dense and/or mature weed infestations. |

Cotton Preplant Application

The maximum preplant application rate is 3 fl oz per acre (0.067 lb ai per acre). Do not exceed a maximum cumulative amount of 6 fl oz per acre per year (0.134 lb ai per acre per year) for all uses on cotton.

| Application Timing | Rate Range (fl oz/A) | Additional Information & Restrictions |
|--------------------|----------------------|--|
| Preplant Burndown | 1 to 3 | <ul style="list-style-type: none">• Apply as a broadcast spray using conventional low-pressure ground spray equipment or by aerial application.• Follow manufacture's recommendations for spraying pressure.• Review and follow restrictions from the spray drift management section.• Do not apply less than 14 days between application and planting.• Use higher rate for dense and/or mature weed infestations.• Do not apply more than once per crop season. |

Soybean

The maximum single application rate is 3 fl oz per acre (0.067 lb ai per acre). Do not exceed 6 fl oz per acre per year (0.134 lb ai per acre per year).

| Application Timing | Rate Range (fl oz/A) | Additional Information & Restrictions |
|--------------------|----------------------|--|
| Preplant Burndown | 1 to 3 | <ul style="list-style-type: none">• Apply as a broadcast spray using conventional low-pressure ground spray equipment or by aerial application.• Follow manufacture's recommendations for spraying pressure.• Review and follow restrictions from the spray drift management section.• Do not reapply within 14 days.• Do not apply less than 14 days between application and planting.• Use higher rate for dense and/or mature weed infestations.• |

NOTICE

Read the "WARRANTY DISCLAIMER" and "LIMITATION OF LIABILITY" in the label booklet for this product before using this product. Those terms apply to this supplemental labeling and if those terms are not acceptable, return the product unopened at once.

(RV021122)

Manufactured for:



HELM Agro US, Inc.
401 E. Jackson St., Suite 1400
Tampa, FL 33602
[Phone: 813-621-8846](tel:813-621-8846)